



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

ed the hypothesis either of right angle or obtuse angle, and from the other the hypothesis of acute angle; which is contrary to P.V, P.VI, and P.VII.

Therefore the three angles together of either of the aforesaid triangles will be less than two right angles; and therefore (P.IX.) is established the hypothesis of acute angle. Quod erat tertio loco demonstrandum.

Accordingly by any triangle  $ABC$ , of which the three angles are equal to, or greater, or less than two right angles, is established respectively the hypothesis of right angle, or obtuse angle, or acute angle. Quod erat propositum.

**COROLLARY.** Hence, any one side of any proposed triangle being produced, as suppose  $AB$  to  $H$ , the external angle  $HBC$  will be (Eu.I.13.) either equal to, or less, or greater than the remaining internal and opposite angles together at the points  $A$  and  $C$ , according as is true the hypothesis of right angle, or obtuse angle, or acute angle. And inversely.

[To be continued.]

---

## THE "IRREDUCIBLE CASE."

---

By J. K. ELLWOOD, A. M., Principal Colfax School, Pittsburg, Pa.

---

**PROBLEM.**—To extract the cube root of  $a \pm \sqrt{-b}$ .

Put  $\sqrt[3]{a + \sqrt{-b}} = m + n$ , and  $\sqrt[3]{a - \sqrt{-b}} = m - n$ .

Then  $a + \sqrt{-b} = m^3 + 3m^2n + 3mn^2 + n^3$ , and  $a - \sqrt{-b} = m^3 - 3m^2n + 3mn^2 - n^3$ . Hence  $a = m^3 + 3mn^2$ , and  $\sqrt{-b} = 3m^2n + n^3$ .

*Example 1.* Find the cube root of  $9 + 25\sqrt{-2}$ .

Here  $a = m^3 + 3mn^2 = 9 = 3^3 - 18$ . Hence  $3mn^2 = -18$ , and  $n = \sqrt{-2}$ .

To verify these values of  $m$  and  $n$  substitute them in  $\sqrt{-b} = 3m^2n + n^3 = 25\sqrt{-2}$ . Doing this we have  $27\sqrt{-2} + (-2\sqrt{-2}) = 25\sqrt{-2}$ .

$\therefore 3 + \sqrt{-2}$  is the required root. When the substituted values of  $m$  and  $n$  do not give the second term they are not correct, and other values must be found by trial.

*Example 2.* Find the cube root of  $2\sqrt{11} + 30\sqrt{-3}$ . Here  $a = m^3 + 3mn^2 = 2\sqrt{11} = (\sqrt{11})^3 - 9\sqrt{11}$ .

Hence  $3mn^2 = -9\sqrt{11}$ , and  $n = \sqrt{-3}$ . Since these values of  $m$  and  $n$  substituted in  $3m^2n + n^3$  give  $\sqrt{-b} = 30\sqrt{-3}$ , the root is  $\sqrt{11} + \sqrt{-3}$ .

This method frequently enables us to simplify Cardan's formula for cubics in what is called the "irreducible case", said formula being

$$x = \sqrt[3]{q + \sqrt{(q^2 - p^3)}} + \sqrt[3]{q - \sqrt{(q^2 - p^3)}}.$$

1. \*  $x^3 - 22x - 24 = 0$ . Here  $p = \frac{2}{3}$ ,  $q = 12$ .

$$\text{Then } x = \sqrt[3]{12 + \frac{2}{3}\sqrt{(-\frac{1}{3})}} + \sqrt[3]{12 - \frac{2}{3}\sqrt{(-\frac{1}{3})}}.$$

Now,  $a = m^3 + 3mn^2 = 12 = -2^3 + 20$ .  $3mn^2 = 20$ ,  $n = \sqrt{-\frac{1}{3}}$ .

$\therefore -2 \pm \sqrt{-\frac{1}{3}}$  are the roots. Then  $x = (-2 + \sqrt{-\frac{1}{3}}) + (-2 - \sqrt{-\frac{1}{3}}) = -4$ .

2.  $x^3 - 8x^2 + 19x - 12 = 0$ . Put  $x = y + \frac{8}{3}$ .

Then  $y^3 - \frac{7}{3}y + \frac{2}{27} = 0$ , where  $p = \frac{7}{3}$ ,  $q = -\frac{1}{27}$ .

$$\begin{aligned} \text{Then } y &= \sqrt[3]{-\frac{1}{27} + \sqrt{(-\frac{7}{3}\frac{2}{27})}} + \sqrt[3]{-\frac{1}{27} - \sqrt{(-\frac{7}{3}\frac{2}{27})}} \\ &= \sqrt[3]{-\frac{1}{27} + \frac{1}{3}\sqrt{-3}} + \sqrt[3]{-\frac{1}{27} - \frac{1}{3}\sqrt{-3}} \end{aligned}$$

Here  $a = m^3 + 3mn^2 = -\frac{1}{27} = (\frac{1}{3})^3 - \frac{1}{27}$ . Hence  $n = \sqrt{-\frac{1}{3}} = \frac{1}{3}\sqrt{-3}$ .

$\therefore y = (\frac{1}{3} + \frac{1}{3}\sqrt{-3}) + (\frac{1}{3} - \frac{1}{3}\sqrt{-3}) = \frac{2}{3}$ . Then  $x = \frac{2}{3} + \frac{8}{3} = 4$ .

3.  $x^3 - 12x^2 + 41x - 42 = 0$ . Put  $x = y + 4$ .

Then  $y^3 - 7y = 6$ , where  $p = \frac{7}{3}$ ,  $q = 3$ . Then  $y = \sqrt[3]{3 + \frac{1}{3}\sqrt{-3}} + \sqrt[3]{3 - \frac{1}{3}\sqrt{-3}}$ .

Here  $a = 3 = (\frac{3}{3})^3 - \frac{3}{3}$ .  $\therefore n = \frac{1}{3}\sqrt{-3}$ .

$\therefore y = (\frac{3}{3} + \frac{1}{3}\sqrt{-3}) + (\frac{3}{3} - \frac{1}{3}\sqrt{-3}) = 3$ , and  $x = 3 + 4 = 7$ .

\* Maynard in his Key to Bonnycastle's Introduction, page 78, says this "can only be resolved by a table of sines, or by infinite series."

## THEOREM 16 OF LOBATSCHESKY'S THEORY OF PARALLELS.

By JOHN N. LYLE, Ph. D., Professor of Natural Sciences, Westminster College, Fulton, Missouri

Says Lobatschewsky in his Theorem 16—"All straight lines which in a plane go out from a point can, with reference to a given straight line in the same plane, be divided into two classes—into *cutting* and *not-cutting*. The *boundary lines* of the one and the other class of those lines will be called *parallel to the given line*.

From the point  $A$  (Fig. 1) let fall upon the line  $BC$  the perpendicular  $AD$ , to which again draw the perpendicular  $AE$ . In the right angle  $EAD$  either will all all straight lines which go out from the point  $A$  meet the line  $DC$ , as for example  $AF$ , or some of them, like the perpendicular  $AE$ , will not meet the line  $DC$ . In the uncertainty whether the perpendicular  $AE$  is the only line which does not meet  $DC$ , we will assume it may be possible that there are still other lines, for




Fig. 1.